


Tarphius furtadoi, Ironclad Beetle

Assessment by: Borges, P.A.V. & Lamelas-López, L.



View on www.iucnredlist.org

Citation: Borges, P.A.V. & Lamelas-López, L. 2018. *Tarphius furtadoi*. The IUCN Red List of Threatened Species 2018: e.T112214889A112214899. <http://dx.doi.org/10.2305/IUCN.UK.2018-1.RLTS.T112214889A112214899.en>

Copyright: © 2018 International Union for Conservation of Nature and Natural Resources

Reproduction of this publication for educational or other non-commercial purposes is authorized without prior written permission from the copyright holder provided the source is fully acknowledged.

Reproduction of this publication for resale, reposting or other commercial purposes is prohibited without prior written permission from the copyright holder. For further details see [Terms of Use](#).

The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#). The IUCN Red List Partners are: [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); and [Zoological Society of London](#).

If you see any errors or have any questions or suggestions on what is shown in this document, please provide us with [feedback](#) so that we can correct or extend the information provided.

Taxonomy

Kingdom	Phylum	Class	Order	Family
Animalia	Arthropoda	Insecta	Coleoptera	Zopheridae

Taxon Name: *Tarphius furtadoi* Borges & Serrano, 2017

Synonym(s):

- *Tarphius azoricus*
- *Tarphius tornvalli*

Common Name(s):

- English: Ironclad Beetle

Taxonomic Source(s):

Borges, P.A.V., Amormin, I.R., Terzopoulou, S. Rigal, F., Emerson, B.C. and Serrano, A.R.M. 2017. Cryptic diversity in the Azorean beetle genus *Tarphius* Erichson, 1845 (Coleoptera: Zopheridae): An integrative taxonomic approach with description of four new species. *Zootaxa* 4236(3): 401-449 DOI: <http://dx.doi.org/10.11646/zootaxa.4236.3.1>.

Taxonomic Notes:

Tarphius tornvalli in Gillerfors (1986, p. 21) (Pico Isl.) *Tarphius tornvalli* in Borges (1990, p. 112) (Pico Isl.) *Tarphius tornvalli* in Borges (1991, p. 2) (Pico Isl.) *Tarphius tornvalli* and *Tarphius azoricus* in Borges *et al.* (2005, p. 207) (Pico Isl.) *Tarphius tornvalli* and *Tarphius azoricus* in Oromí *et al.* (2010, p. 232) (Pico Isl.) *Tarphius tornvalli* and *Tarphius azoricus* in Amorim *et al.* (2012, Fig. 2) (Pico Isl.) *Tarphius azoricus* in Borges *et al.* (2005, p. 207) (São Jorge Isl.) *Tarphius azoricus* in Oromí *et al.* (2010, p. 232) (São Jorge Isl.) *Tarphius azoricus* in Amorim *et al.* (2012, Fig. 2) (São Jorge Isl.) *Tarphius azoricus* in Oromí *et al.* (2010, p. 232) (Faial Isl.) *Tarphius azoricus* in Amorim *et al.* (2012, Fig. 2) (Faial Isl.)

Assessment Information

Red List Category & Criteria: Endangered B1ab(i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v) [ver 3.1](#)

Year Published: 2018

Date Assessed: April 5, 2017

Justification:

Tarphius furtadoi is an endemic species present in S. Jorge, Pico and Faial islands (Azores, Portugal) (Borges *et al.* 2017). It has an extent of occurrence of 890 km² and an area of occupancy of 48 km². There is a continuing decline in the EOO, AOO, extent and quality of habitat as well as the number of mature individuals as a result of the invasions of non-native plants. The species occurs mainly in the soil of native forests, but also under the bark and dead wood of endemic and exotic trees (e.g. *Cryptomeria japonica* and *Acacia* sp.). In the past, the species has probably strongly declined due to changes in habitat size. Therefore, we suggest as future measures of conservation: (1) a long-term monitoring plan of the species; and (2) control of invasive species, namely *Hedychium gardnerianum*. The species is assessed as Endangered (EN).

Geographic Range

Range Description:

Tarphius furtadoi is an endemic species present in S. Jorge, Pico and Faial islands (Azores, Portugal) (Borges *et al.* 2017), known from Natural Reserves of Caveiro, Lagoa do Caiado and Mistério da Prainha (Pico island); Pico Pinheiro and Topo (S. Jorge island) and Cabeço do Fogo and Caldeira do Faial (Faial island). The extent of occurrence (EOO) is *ca* 890 km² and the maximum estimated area of occupancy (AOO) is 48 km².

Country Occurrence:

Native: Portugal (Azores)

Distribution Map

Tarphius furtadoi




Range

Extant (resident)

Compiled by:

Paulo Borges



The boundaries and names shown and the designations used on this map do not imply any official endorsement, acceptance or opinion by IUCN.



Population

The species is abundant, particularly in the well preserved patches of native forests of S. Jorge, Pico and Faial islands (Borges *et al.* 2017). A continuing decline in the number of mature individuals is inferred from monitoring schemes and from the ongoing habitat degradation due to invasions of alien plants (Borges *et al.* 2017). This species is assessed here as severely fragmented as at least 50% of its population can be found in subpopulations that are 1) smaller than would be required to support a viable population, and 2) separated from other habitat patches by a large distance. In fact, the species occurs in fragments that are isolated in a matrix of pastures.

Current Population Trend: Decreasing

Habitat and Ecology (see Appendix for additional information)

The species is particularly abundant, namely this species live in the soil and occurring in some of the larger and well preserved patches of native forests of S. Jorge, Pico and Faial islands. It also occurs under the bark of dead wood of endemic and exotic trees (Borges *et al.* 2017). This species has an altitudinal range between 250 and 1000 m. It is a nocturnal fungivorous species.

Systems: Terrestrial

Use and Trade

This species is not utilised.

Threats (see Appendix for additional information)

In the past, the species has probably strongly declined due to changes in habitat size and quality (Triantis *et al.* 2010). Currently, the rapid advance and expansion of invasive plants species is the major threat (Borges *et al.* 2017), particularly *Hedychium gardnerianum* and *Pittosporum undulatum* since are changing the habitat structure, namely decreasing the cover of bryophytes and ferns in the soil and promoting the spread of other plants. The management of *Cryptomeria japonica* plantations could be also a problem for the subpopulations living in this habitat. Based on Ferreira *et al.* 2016 the habitat will further decline as a consequence of climate change (increasing number of droughts and habitat shifting and alteration). The future removal of some patches with exotic vegetation for pasture implementation may also be a threat for some subpopulations occurring in this marginal habitat.

Conservation Actions (see Appendix for additional information)

The species is not protected by regional law. Its habitat is in regionally protected areas (Natural Forest Reserves of Mistério da Prainha, Caveiro and Caiado in Pico island; Pico Pinheiro and Topo in S. Jorge island and Cabeço do Fogo and Caldeira do Faial in Faial island). Degraded habitats should be restored with the removal of invasive species. A strategy needs also to be developed to address the future threat by climate change. A habitat management plan is needed and anticipated to be developed during the coming years. Since this species is an icon of the relict native Azorean forests, it is suggested that some awareness measures should be put in practice. Further research is needed into its ecology and life history in order to find extant specimens in additional native forest fragments and obtain information on population size, distribution and trends. It is also necessary an area-based management plan and a monitoring plan for the invertebrate community in the habitat in order to contribute to perform a

species potential recovery plan. A monitoring every ten years using the BALA protocol will inform about habitat quality (see e.g. Gaspar *et al.* 2011).

Credits

Assessor(s): Borges, P.A.V. & Lamelas-López, L.

Reviewer(s): Danielczak, A.

Bibliography

Amorim, I., Emerson, B.C, Borges, P.A.V. & Wayne, R.K. 2012. Phylogeography and molecular phylogeny of Macaronesian island *Tarphius*. *Journal of Biogeography* 39: 1583–1595.

Borges, P.A.V. 1990. A checklist of Coleoptera from the Azores with some systematic and biogeographic comments. *Boletim do Museu Municipal do Funchal* 42(87-136).

Borges, P.A.V. 1991. Two new species of *Tarphius* Erichson, 1848 (Coleoptera, Colydiidae) from the Azores. *Bocagiana* 143: 1-11.

Borges, P.A.V., Amormin, I.R., Terzopoulou, S. Rigal, F., Emerson, B.C. and Serrano, A.R.M. 2017. Cryptic diversity in the Azorean beetle genus *Tarphius* Erichson, 1845 (Coleoptera: Zopheridae): An integrative taxonomic approach with description of four new species. *Zootaxa* 4236(3): 401-449 DOI: <http://dx.doi.org/10.11646/zootaxa.4236.3.1>.

Borges, P.A.V., Costa, A., Cunha, R., Gabriel, R., Gonçalves, V., Martins, A.F., Melo, I., Parente, M., Raposeiro, P., Rodrigues, P., Santos, R.S., Silva, L., Vieira, P. & Vieira, V. 2010. *A list of the terrestrial and marine biota from the Azores*. Princípiã, Cascais.

Borges, P.A.V., Cunha, R., Gabriel, R., Martins, A.F., Silva, L. and Vieira, V. (eds). 2005. *A list of the terrestrial fauna (Mollusca and Arthropoda) and flora (Bryophyta, Pteridophyta and Spermatophyta) from the Azores*. pp. 318. Horta, Angra do Heroísmo and Ponta Delgada.

Ferreira, M.T., Cardoso, P., Borges, P.A.V., Gabriel, R., Azevedo, E.B., Reis, F., Araújo, M.B. and Elias, R.B. 2016. Effects of climate change on the distribution of indigenous species in oceanic islands (Azores). *Climate Change* 138: 603-615.

Gillerfors, G. 1986. Contribution to the coleopterous fauna of the Azores. *Boletim do Museu Municipal do Funchal* 38(172): 16-27.

IUCN. 2018. The IUCN Red List of Threatened Species. Version 2018-1. Available at: www.iucnredlist.org. (Accessed: 28 June 2018).

Triantis, K.A., Borges, P.A.V., Ladle, R.J., Hortal, J., Cardoso, P., Gaspar, C., Dinis, F., Mendonça, E., Silveira, L.M.A., Gabriel, R., Melo, C., Santos, A.M.C., Amorim, I.R., Ribeiro, S.P., Serrano, A.R.M., Quartau, J.A. and Whittaker, R.J. 2010. Extinction debt on oceanic islands. *Ecography* 33: 285-294.

Citation

Borges, P.A.V. & Lamelas-López, L. 2018. *Tarphius furtadoi*. The IUCN Red List of Threatened Species 2018: e.T112214889A112214899. <http://dx.doi.org/10.2305/IUCN.UK.2018-1.RLTS.T112214889A112214899.en>

Disclaimer

To make use of this information, please check the [Terms of Use](#).

External Resources

For [Images and External Links to Additional Information](#), please see the [Red List website](#).

Appendix

Habitats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Habitat	Season	Suitability	Major Importance?
1. Forest -> 1.3. Forest - Subantarctic	Resident	Suitable	Yes
14. Artificial/Terrestrial -> 14.3. Artificial/Terrestrial - Plantations	Resident	Suitable	No
0. Root -> 16. Introduced vegetation	Resident	Suitable	Yes

Threats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Threat	Timing	Scope	Severity	Impact Score
11. Climate change & severe weather -> 11.1. Habitat shifting & alteration	Future	Whole (>90%)	Slow, significant declines	Low impact: 5
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality 2. Species Stresses -> 2.2. Species disturbance		
11. Climate change & severe weather -> 11.2. Droughts	Ongoing	Whole (>90%)	Slow, significant declines	Medium impact: 7
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality		
2. Agriculture & aquaculture -> 2.2. Wood & pulp plantations -> 2.2.1. Small-holder plantations	Ongoing	Minority (50%)	Causing/could cause fluctuations	Low impact: 5
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality 2. Species Stresses -> 2.2. Species disturbance		
2. Agriculture & aquaculture -> 2.3. Livestock farming & ranching -> 2.3.2. Small-holder grazing, ranching or farming	Future	Minority (50%)	Very rapid declines	Low impact: 5
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation 1. Ecosystem stresses -> 1.3. Indirect ecosystem effects 2. Species Stresses -> 2.1. Species mortality 2. Species Stresses -> 2.2. Species disturbance		
8. Invasive and other problematic species, genes & diseases -> 8.1. Invasive non-native/alien species/diseases -> 8.1.2. Named species (Hedychium gardnerianum)	Ongoing	Majority (50-90%)	Rapid declines	Medium impact: 7
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.2. Species disturbance		

8. Invasive and other problematic species, genes & diseases -> 8.1. Invasive non-native/alien species/diseases -> 8.1.2. Named species (Pittosporum undulatum)	Ongoing	Majority (50-90%)	Slow, significant declines	Medium impact: 6
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.2. Species disturbance		

Conservation Actions in Place

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Conservation Actions in Place
In-Place Land/Water Protection and Management
Conservation sites identified: Yes, over part of range
Occur in at least one PA: Yes
Percentage of population protected by PAs (0-100): 71-80

Conservation Actions Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Conservation Actions Needed
2. Land/water management -> 2.1. Site/area management
2. Land/water management -> 2.2. Invasive/problematic species control
2. Land/water management -> 2.3. Habitat & natural process restoration
4. Education & awareness -> 4.1. Formal education
4. Education & awareness -> 4.3. Awareness & communications
5. Law & policy -> 5.4. Compliance and enforcement -> 5.4.3. Sub-national level

Research Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Research Needed
1. Research -> 1.2. Population size, distribution & trends
1. Research -> 1.3. Life history & ecology
2. Conservation Planning -> 2.2. Area-based Management Plan
3. Monitoring -> 3.1. Population trends
3. Monitoring -> 3.4. Habitat trends

Additional Data Fields

Distribution
Estimated area of occupancy (AOO) (km ²): 48
Continuing decline in area of occupancy (AOO): Yes
Extreme fluctuations in area of occupancy (AOO): Unknown
Estimated extent of occurrence (EOO) (km ²): 890
Continuing decline in extent of occurrence (EOO): Yes
Extreme fluctuations in extent of occurrence (EOO): Unknown
Number of Locations: 8
Continuing decline in number of locations: Yes
Extreme fluctuations in the number of locations: Unknown
Lower elevation limit (m): 250
Upper elevation limit (m): 1000
Population
Continuing decline of mature individuals: Yes
Extreme fluctuations: Unknown
Population severely fragmented: Yes
Habitats and Ecology
Continuing decline in area, extent and/or quality of habitat: Yes
Generation Length (years): 1
Movement patterns: Not a Migrant

The IUCN Red List Partnership



The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#).

The IUCN Red List Partners are: [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); and [Zoological Society of London](#).