


Agabus godmanni, Azorean predacious diving beetle

Assessment by: Borges, P.A.V. & Lamelas-López, L.



View on www.iucnredlist.org

Citation: Borges, P.A.V. & Lamelas-López, L. 2018. *Agabus godmanni*. The IUCN Red List of Threatened Species 2018: e.T97163447A99166669. <http://dx.doi.org/10.2305/IUCN.UK.2018-1.RLTS.T97163447A99166669.en>

Copyright: © 2018 International Union for Conservation of Nature and Natural Resources

Reproduction of this publication for educational or other non-commercial purposes is authorized without prior written permission from the copyright holder provided the source is fully acknowledged.

Reproduction of this publication for resale, reposting or other commercial purposes is prohibited without prior written permission from the copyright holder. For further details see [Terms of Use](#).

The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#). The IUCN Red List Partners are: [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); and [Zoological Society of London](#).

If you see any errors or have any questions or suggestions on what is shown in this document, please provide us with [feedback](#) so that we can correct or extend the information provided.

Taxonomy

Kingdom	Phylum	Class	Order	Family
Animalia	Arthropoda	Insecta	Coleoptera	Dytiscidae

Taxon Name: *Agabus godmanni* Crotch, 1867

Common Name(s):

- English: Azorean predacious diving beetle

Taxonomic Source(s):

2016. Integrated Taxonomic Information System. Available at: <http://www.itis.gov/>.

Assessment Information

Red List Category & Criteria: Endangered B2ab(i,ii,iii,iv,v) [ver 3.1](#)

Year Published: 2018

Date Assessed: January 3, 2017

Justification:

Agabus godmani is an endemic species present in Flores, Faial, S. Jorge, Terceira and S. Miguel islands (Azores, Portugal). It has a large extent of occurrence (EOO = ca 17,600 km²) and small area of occupancy (AOO = 136 km²). The species is common and known from at least five fragmented subpopulations in 10 locations. In the past, the species has probably strongly declined due to changes in habitat size and quality. The main threat to this species is the impact of introduced species and the agricultural activities (landscape transformation, pollution, habitat destruction). Based on Ferreira *et al.* (2016) the habitat will further decline as a consequence of climate change (increasing number of droughts). Based upon the small area of occupancy and associated decline in extent of occurrence, area of occupancy and habitat quality it is assessed as Endangered.

Geographic Range

Range Description:


Agabus godmani is an endemic species present in Flores, Faial, S. Jorge, Terceira and S. Miguel islands (Azores, Portugal) (Borges *et al.* 2010), known from Natural Forest Reserves of Topo (S. Jorge); Morro Alto e Pico da Sé (Flores), Caldeira do Faial (Faial), Graminhais and Pico da Vara (S. Miguel). The extent of occurrence (EOO) is ca 17,600 km² and the maximum estimated area of occupancy (AOO) is 136 km².

Country Occurrence:

Native: Portugal (Azores)

Distribution Map

Agabus godmanni



Range

Extant (resident)

Compiled by:

Paulo Borges



The boundaries and names shown and the designations used on this map do not imply any official endorsement, acceptance or opinion by IUCN.



Population

The species is rare in all known subpopulations in freshwater areas in several islands (Flores, Faial, S. Jorge, Terceira and S. Miguel islands). A continuing decline in the number of mature individuals is inferred from the ongoing habitat degradation due to human activities (habitat transformation associated to agricultural activities and cattle pollution) and introduced species. This species is assessed as severely fragmented, as it is distributed in isolated freshwater areas in several islands. At least 50% of its population can be found in subpopulations/in habitat patches that are 1) smaller than would be required to support a viable population, and 2) separated from other habitat patches by a large distance. In fact, the species occurs in fragments that are isolated in a sea of pastures and *Cryptomeria japonica* plantations.

Current Population Trend: Decreasing

Habitat and Ecology (see Appendix for additional information)

The species occurs in the freshwater habitats located in native forests of the Flores, Faial, S. Jorge, Terceira and S. Miguel islands (Azores), with an altitudinal range between 450 and 1200 m. Adults and larvae are predators that live in the water column of freshwater systems.

Systems: Freshwater

Use and Trade

The species is not utilised.

Threats (see Appendix for additional information)

The main threat to this species is the impact of introduced species and the agricultural activities (landscape transformation, pollution by herbicides and pesticides and nutrient loads, habitat destruction). However, we may add to these the impact of climatic changes with major impact on habitat shifting & alteration and droughts (Ferreira *et al.* 2016).

Conservation Actions (see Appendix for additional information)

The species is not protected by regional law. Its habitat is in regionally protected areas (Natural Parks of Flores, Faial, S. Jorge and S. Miguel). Degraded habitats should be restored and a strategy needs to be developed to address the future threat by climate change. Further research is needed into its ecology and life history in order to find extant specimens. It is necessary a monitoring plan for the invertebrate community in the habitat in order to contribute to the conservation of this species. A habitat management plan is needed and anticipated to be developed during the coming years. Monitoring every ten years using the BALA protocol will inform about habitat quality (see e.g. Gaspar *et al.* 2010).

Credits

Assessor(s): Borges, P.A.V. & Lamelas-López, L.

Reviewer(s): Danielczak, A.

Bibliography

Borges, P.A.V., Costa, A., Cunha, R., Gabriel, R., Gonçalves, V., Martins, A.F., Melo, I., Parente, M., Raposeiro, P., Rodrigues, P., Santos, R.S., Silva, L., Vieira, P. & Vieira, V. 2010. *A list of the terrestrial and marine biota from the Azores*. Princípiã, Cascais.

Ferreira, M.T., Cardoso, P., Borges, P.A.V., Gabriel, R., Azevedo, E.B., Reis, F., Araújo, M.B. and Elias, R.B. 2016. Effects of climate change on the distribution of indigenous species in oceanic islands (Azores). *Climate Change* 138: 603-615.

Gaspar, C., Gaston, K.J., Borges, P.A.V. and Cardoso, P. 2011. Selection of priority areas for arthropod conservation in the Azores archipelago. *Journal of Insect Conservation* 15: 671–684.

IUCN. 2018. The IUCN Red List of Threatened Species. Version 2018-1. Available at: www.iucnredlist.org. (Accessed: 28 June 2018).

Citation

Borges, P.A.V. & Lamelas-López, L. 2018. *Agabus godmanni*. The IUCN Red List of Threatened Species 2018: e.T97163447A99166669. <http://dx.doi.org/10.2305/IUCN.UK.2018-1.RLTS.T97163447A99166669.en>

Disclaimer

To make use of this information, please check the [Terms of Use](#).

External Resources

For [Images and External Links to Additional Information](#), please see the [Red List website](#).

Appendix

Habitats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Habitat	Season	Suitability	Major Importance?
1. Forest -> 1.4. Forest - Temperate	Resident	Suitable	Yes
5. Wetlands (inland) -> 5.1. Wetlands (inland) - Permanent Rivers/Streams/Creeks (includes waterfalls)	Resident	Suitable	Yes
5. Wetlands (inland) -> 5.2. Wetlands (inland) - Seasonal/Intermittent/Irregular Rivers/Streams/Creeks	Resident	Suitable	Yes
5. Wetlands (inland) -> 5.7. Wetlands (inland) - Permanent Freshwater Marshes/Pools (under 8ha)	Resident	Suitable	Yes
15. Artificial/Aquatic & Marine -> 15.2. Artificial/Aquatic - Ponds (below 8ha)	Resident	Suitable	Yes

Threats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Threat	Timing	Scope	Severity	Impact Score
10. Geological events -> 10.1. Volcanoes	Future	Whole (>90%)	Very rapid declines	Medium impact: 7
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 2. Species Stresses -> 2.1. Species mortality		
11. Climate change & severe weather -> 11.1. Habitat shifting & alteration	Future	Whole (>90%)	Slow, significant declines	Low impact: 5
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality 2. Species Stresses -> 2.2. Species disturbance		
11. Climate change & severe weather -> 11.2. Droughts	Ongoing	Whole (>90%)	Slow, significant declines	Medium impact: 7
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality		
2. Agriculture & aquaculture -> 2.2. Wood & pulp plantations -> 2.2.1. Small-holder plantations	Ongoing	Minority (50%)	Causing/could cause fluctuations	Low impact: 5
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality 2. Species Stresses -> 2.2. Species disturbance		
2. Agriculture & aquaculture -> 2.3. Livestock farming & ranching -> 2.3.2. Small-holder grazing, ranching or farming	Ongoing	Minority (50%)	Slow, significant declines	Low impact: 5
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation		

			2. Species Stresses -> 2.1. Species mortality	
			2. Species Stresses -> 2.2. Species disturbance	
8. Invasive and other problematic species, genes & diseases -> 8.1. Invasive non-native/alien species/diseases -> 8.1.1. Unspecified species	Ongoing	Majority (50-90%)	Slow, significant declines	Medium impact: 6
	Stresses:		1. Ecosystem stresses -> 1.2. Ecosystem degradation	
			2. Species Stresses -> 2.1. Species mortality	
			2. Species Stresses -> 2.2. Species disturbance	
9. Pollution -> 9.3. Agricultural & forestry effluents -> 9.3.1. Nutrient loads	Ongoing	Majority (50-90%)	Slow, significant declines	Medium impact: 6
	Stresses:		1. Ecosystem stresses -> 1.2. Ecosystem degradation	
			2. Species Stresses -> 2.1. Species mortality	
			2. Species Stresses -> 2.2. Species disturbance	
9. Pollution -> 9.3. Agricultural & forestry effluents -> 9.3.3. Herbicides and pesticides	Ongoing	Majority (50-90%)	Rapid declines	Medium impact: 7
	Stresses:		1. Ecosystem stresses -> 1.2. Ecosystem degradation	
			2. Species Stresses -> 2.1. Species mortality	
			2. Species Stresses -> 2.2. Species disturbance	

Conservation Actions in Place

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Conservation Actions in Place
In-Place Land/Water Protection and Management
Conservation sites identified: Yes, over part of range
Occur in at least one PA: Yes
Percentage of population protected by PAs (0-100): 81-90

Conservation Actions Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Conservation Actions Needed
2. Land/water management -> 2.1. Site/area management
2. Land/water management -> 2.2. Invasive/problematic species control
2. Land/water management -> 2.3. Habitat & natural process restoration
4. Education & awareness -> 4.1. Formal education
4. Education & awareness -> 4.3. Awareness & communications
5. Law & policy -> 5.4. Compliance and enforcement -> 5.4.3. Sub-national level

Research Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Research Needed
1. Research -> 1.2. Population size, distribution & trends
1. Research -> 1.3. Life history & ecology
1. Research -> 1.5. Threats
2. Conservation Planning -> 2.2. Area-based Management Plan
3. Monitoring -> 3.1. Population trends
3. Monitoring -> 3.4. Habitat trends

Additional Data Fields

Distribution
Estimated area of occupancy (AOO) (km ²): 153
Continuing decline in area of occupancy (AOO): Yes
Extreme fluctuations in area of occupancy (AOO): Unknown
Estimated extent of occurrence (EOO) (km ²): 17600
Continuing decline in extent of occurrence (EOO): Yes
Extreme fluctuations in extent of occurrence (EOO): Unknown
Number of Locations: 10
Continuing decline in number of locations: Yes
Extreme fluctuations in the number of locations: Unknown
Lower elevation limit (m): 450
Upper elevation limit (m): 1200
Population
Continuing decline of mature individuals: Yes
Population severely fragmented: Yes
Habitats and Ecology
Continuing decline in area, extent and/or quality of habitat: Yes
Generation Length (years): 1
Movement patterns: Not a Migrant

The IUCN Red List Partnership



The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#).

The IUCN Red List Partners are: [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); and [Zoological Society of London](#).