


Neomariania scriptella, Moth

Assessment by: Vieira, V. & Borges, P.A.V.



View on www.iucnredlist.org

Citation: Vieira, V. & Borges, P.A.V. 2018. *Neomariania scriptella*. The IUCN Red List of Threatened Species 2018: e.T97239466A99166949. <http://dx.doi.org/10.2305/IUCN.UK.2018-1.RLTS.T97239466A99166949.en>

Copyright: © 2018 International Union for Conservation of Nature and Natural Resources

Reproduction of this publication for educational or other non-commercial purposes is authorized without prior written permission from the copyright holder provided the source is fully acknowledged.

Reproduction of this publication for resale, reposting or other commercial purposes is prohibited without prior written permission from the copyright holder. For further details see [Terms of Use](#).

The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#). The IUCN Red List Partners are: [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); and [Zoological Society of London](#).

If you see any errors or have any questions or suggestions on what is shown in this document, please provide us with [feedback](#) so that we can correct or extend the information provided.

Taxonomy

Kingdom	Phylum	Class	Order	Family
Animalia	Arthropoda	Insecta	Lepidoptera	Stathmopodidae

Taxon Name: *Neomariania scriptella* Rebel, 1940

Synonym(s):

- *Megaceraea scriptella* Rebel, 1940

Common Name(s):

- English: Moth

Taxonomic Source(s):

De Jong, Y., Verbeek, M., Michelsen, V., Bjørn, P.P., Los, W., Steeman, F., Bailly, N., Basire, C., Chylarecki, P., Stloukal, E., Hagedorn, G., Wetzels, F.T., Glöckler, F., Kroupa, A., Korb, G., Hoffmann, A., Häuser, C., Kohlbecker, A., Müller, A., Güntsch, A., Stoev, P. and Penev, L. 2014. Fauna Europaea – all European animal species on the web. *Biodiversity Data Journal* 2: e4034. DOI: 10.3897/BDJ.2.e4034.

Assessment Information

Red List Category & Criteria: Endangered B1ab(i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v) [ver 3.1](#)

Year Published: 2018

Date Assessed: March 24, 2017

Justification:

Neomariania scriptella is an endemic species present in Pico, Graciosa and Terceira islands (Azores, Portugal) (Borges *et al.* 2010). It has a relatively large extent of occurrence (EOO = ca 3,400 km²), but a small area of occupancy (AOO = 48 km²). Currently *Neomariania scriptella* is under threat due to degradation of the habitat by urban development and agriculture, but also due to invasive plants *Pittosporum undulatum* and *Hedychium gardnerianum* that are changing some of the areas and decreasing the quality of the habitat. Based on Ferreira *et al.* (2016) the habitat will further decline as a consequence of climate change. Based upon the small area of occupancy, decreasing quality of the habitat and low number of subpopulations it is assessed as Endangered.

Geographic Range

Range Description:

Neomariania scriptella is an endemic species present in Pico, Graciosa and Terceira islands (Azores, Portugal) (Borges *et al.* 2010), known from native forest and in open landscapes with isolated tree groups. The extent of occurrence (EOO) is ca 3,400 km² and the maximum estimated area of occupancy (AOO) is 48 km².

Country Occurrence:

Native: Portugal (Azores)

Distribution Map

Neomariania scriptella



Range
■ Extant (resident)

Compiled by:
Paulo Borges



Population

Neomariania scriptella is low abundant in three Central Azorean islands (Pico, Graciosa and Terceira), occurring mostly in highly modified native vegetation but also in naturalised plants.

Current Population Trend: Unknown

Habitat and Ecology (see Appendix for additional information)

The species was originally associated with native forest areas but also in open landscapes with isolated tree groups. The vegetation in those sites is currently highly modified. Possibly, the larvae feed on fruit and flowers such as Fabaceae and Moraceae, also in sporangia of ferns; the adult flies in June (Rebel 1940). Altitudinal range: 10-500 m.

Systems: Terrestrial

Use and Trade

This species is not utilised.

Threats (see Appendix for additional information)

In the past, the species has probably strongly declined due to changes in habitat size and quality, mostly the creation of pastures (Triantis *et al.* 2010). Currently several threats are operating: i) urban development; ii) agriculture; iii) invasive plants *Pittosporum undulatum* and *Hedychium gardnerianum* are changing some of the areas and decreasing the quality of the habitat. These changes are decreasing the relative cover of endemic plants and changing the soil cover (decreasing the cover of bryophytes and ferns). Based on Ferreira *et al.* (2016) the habitat will further decline as a consequence of climate change (increasing number of droughts and habitat shifting & alteration).

Conservation Actions (see Appendix for additional information)

The species is not protected by regional law. Its habitat is in regionally protected areas (Natural Park of Graciosa). Further research is needed to monitor the species and conservation measures to control the invasive *Hedychium gardnerianum* and *Pittosporum undulatum* should be implemented to improve habitat quality for this species. Additional research is needed into its ecology and life history in order to find extant specimens. Degraded habitats should be restored and a strategy needs to be developed to address the future threat by climate change. It is necessary a monitoring plan for the invertebrate community in the habitat in order to contribute to the conservation of this species. A habitat management plan is needed and anticipated to be developed during the coming years. Monitoring every ten years using the BALA protocol will inform about habitat quality (see e.g. Gaspar *et al.* 2010).

Credits

Assessor(s): Vieira, V. & Borges, P.A.V.

Reviewer(s): Danielczak, A.

Contributor(s): Nunes, R.

Bibliography

Borges, P.A.V., Costa, A., Cunha, R., Gabriel, R., Gonçalves, V., Martins, A.F., Melo, I., Parente, M., Raposeiro, P., Rodrigues, P., Santos, R.S., Silva, L., Vieira, P. & Vieira, V. 2010. *A list of the terrestrial and marine biota from the Azores*. Princípiã, Cascais.

Ferreira, M.T., Cardoso, P., Borges, P.A.V., Gabriel, R., Azevedo, E.B., Reis, F., Araújo, M.B. and Elias, R.B. 2016. Effects of climate change on the distribution of indigenous species in oceanic islands (Azores). *Climate Change* 138: 603-615.

Gaspar, C., Gaston, K.J., Borges, P.A.V. and Cardoso, P. 2011. Selection of priority areas for arthropod conservation in the Azores archipelago. *Journal of Insect Conservation* 15: 671–684.

IUCN. 2018. The IUCN Red List of Threatened Species. Version 2018-1. Available at: www.iucnredlist.org. (Accessed: 28 June 2018).

Rebel, H. 1940. Die Lepidopterenfauna des Azorischen Archipels. *Commentationes Biologicae* 8(1): 1-59.

Triantis, K.A., Borges, P.A.V., Ladle, R.J., Hortal, J., Cardoso, P., Gaspar, C., Dinis, F., Mendonça, E., Silveira, L.M.A., Gabriel, R., Melo, C., Santos, A.M.C., Amorim, I.R., Ribeiro, S.P., Serrano, A.R.M., Quartau, J.A. and Whittaker, R.J. 2010. Extinction debt on oceanic islands. *Ecography* 33: 285-294.

Citation

Vieira, V. & Borges, P.A.V. 2018. *Neomariania scriptella*. The IUCN Red List of Threatened Species 2018: e.T97239466A99166949. <http://dx.doi.org/10.2305/IUCN.UK.2018-1.RLTS.T97239466A99166949.en>

Disclaimer

To make use of this information, please check the [Terms of Use](#).

External Resources

For [Images and External Links to Additional Information](#), please see the [Red List website](#).

Appendix

Habitats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Habitat	Season	Suitability	Major Importance?
1. Forest -> 1.4. Forest - Temperate	Resident	Suitable	Yes
3. Shrubland -> 3.4. Shrubland - Temperate	Resident	Suitable	Yes
4. Grassland -> 4.4. Grassland - Temperate	Resident	Suitable	Yes
0. Root -> 16. Introduced vegetation	Resident	Marginal	-

Threats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Threat	Timing	Scope	Severity	Impact Score
10. Geological events -> 10.1. Volcanoes	Future	Whole (>90%)	Very rapid declines	Medium impact: 7
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 2. Species Stresses -> 2.1. Species mortality		
1. Residential & commercial development -> 1.1. Housing & urban areas	Ongoing	Minority (50%)	Slow, significant declines	Low impact: 5
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality 2. Species Stresses -> 2.2. Species disturbance		
11. Climate change & severe weather -> 11.1. Habitat shifting & alteration	Future	Whole (>90%)	Slow, significant declines	Low impact: 5
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality 2. Species Stresses -> 2.2. Species disturbance		
11. Climate change & severe weather -> 11.2. Droughts	Ongoing	Whole (>90%)	Slow, significant declines	Medium impact: 7
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality		
2. Agriculture & aquaculture -> 2.1. Annual & perennial non-timber crops -> 2.1.2. Small-holder farming	Ongoing	Minority (50%)	Causing/could cause fluctuations	Low impact: 5
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality 2. Species Stresses -> 2.2. Species disturbance		
2. Agriculture & aquaculture -> 2.3. Livestock farming & ranching -> 2.3.2. Small-holder grazing, ranching or farming	Ongoing	Minority (50%)	Causing/could cause fluctuations	Low impact: 5

	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.1. Species mortality 2. Species Stresses -> 2.2. Species disturbance		
8. Invasive and other problematic species, genes & diseases -> 8.1. Invasive non-native/alien species/diseases -> 8.1.2. Named species (Pittosporum undulatum)	Ongoing	Majority (50-90%)	Rapid declines	Medium impact: 7
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.2. Species disturbance		
8. Invasive and other problematic species, genes & diseases -> 8.1. Invasive non-native/alien species/diseases -> 8.1.2. Named species (Hedychium gardnerianum)	Ongoing	Majority (50-90%)	Rapid declines	Medium impact: 7
	Stresses:	1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.2. Species disturbance		

Conservation Actions in Place

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Conservation Actions in Place
In-Place Research, Monitoring and Planning
Systematic monitoring scheme: Yes
In-Place Land/Water Protection and Management
Conservation sites identified: Yes, over part of range
Occur in at least one PA: Yes
Percentage of population protected by PAs (0-100): 61-70

Conservation Actions Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Conservation Actions Needed
2. Land/water management -> 2.1. Site/area management
2. Land/water management -> 2.2. Invasive/problematic species control
2. Land/water management -> 2.3. Habitat & natural process restoration
4. Education & awareness -> 4.1. Formal education
4. Education & awareness -> 4.3. Awareness & communications
5. Law & policy -> 5.4. Compliance and enforcement -> 5.4.3. Sub-national level

Research Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Research Needed
1. Research -> 1.2. Population size, distribution & trends
1. Research -> 1.3. Life history & ecology
3. Monitoring -> 3.1. Population trends
3. Monitoring -> 3.4. Habitat trends

Additional Data Fields

Distribution
Estimated area of occupancy (AOO) (km ²): 48
Continuing decline in area of occupancy (AOO): Yes
Extreme fluctuations in area of occupancy (AOO): Unknown
Estimated extent of occurrence (EOO) (km ²): 3394
Continuing decline in extent of occurrence (EOO): Yes
Extreme fluctuations in extent of occurrence (EOO): Unknown
Number of Locations: 3
Continuing decline in number of locations: Yes
Extreme fluctuations in the number of locations: Unknown
Lower elevation limit (m): 10
Upper elevation limit (m): 500
Population
Continuing decline of mature individuals: Yes
Population severely fragmented: No
Habitats and Ecology
Continuing decline in area, extent and/or quality of habitat: Yes
Generation Length (years): 1
Movement patterns: Not a Migrant

The IUCN Red List Partnership



The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#).

The IUCN Red List Partners are: [Arizona State University](#); [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); and [Zoological Society of London](#).